



# Non covered vertices in Fibonacci cubes by a maximum set of disjoint hypercubes

Michel Mollard

## ► To cite this version:

Michel Mollard. Non covered vertices in Fibonacci cubes by a maximum set of disjoint hypercubes. Discrete Applied Mathematics, 2017, 219, pp.219-221. hal-01324319v2

**HAL Id: hal-01324319**

**<https://hal.science/hal-01324319v2>**

Submitted on 18 Oct 2016

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Non covered vertices in Fibonacci cubes by a maximum set of disjoint hypercubes

Michel Mollard\*

October 18, 2016

## Abstract

The *Fibonacci cube* of dimension  $n$ , denoted as  $\Gamma_n$ , is the subgraph of  $n$ -cube  $Q_n$  induced by vertices with no consecutive 1's. In this short note we give an immediate proof that asymptotically all vertices of  $\Gamma_n$  are covered by a maximum set of disjoint subgraphs isomorphic to  $Q_k$ , answering an open problem proposed in [2] and solved with a longer proof in [3].

**Keywords:** Fibonacci cube, Fibonacci numbers.

**AMS Subj. Class. (2010):**

## 1 Introduction

Let  $n$  be a positive integer and denote  $[n] = \{1, \dots, n\}$ , and  $[n]_0 = \{0, \dots, n-1\}$ . The  $n$ -cube, denoted as  $Q_n$ , is the graph with vertex set

$$V(Q_n) = \{x_1x_2 \dots x_n \mid x_i \in [2]_0 \text{ for } i \in [n]\},$$

where two vertices are adjacent in  $Q_n$  if the corresponding strings differ in exactly one position. The *Fibonacci  $n$ -cube*, denoted by  $\Gamma_n$ , is the subgraph of  $Q_n$  induced by vertices with no consecutive 1's. Let  $\{F_n\}$  be the *Fibonacci numbers*:  $F_0 = 0$ ,  $F_1 = 1$ ,  $F_n = F_{n-1} + F_{n-2}$  for  $n \geq 2$ . The number of vertices of  $\Gamma_n$  is  $|V(\Gamma_n)| = F_{n+2}$ . Fibonacci cubes have been investigated from many points of view and we refer to the survey [1] for more information about them. Let  $q_k(n)$  be the maximum number of disjoint subgraphs isomorphic to  $Q_k$  in  $\Gamma_n$ . This number is studied in a recent paper [2]. The authors obtained the following recursive formula

**Theorem 1.1** *For every  $k \geq 1$  and  $n \geq 3$   $q_k(n) = q_{k-1}(n-2) + q_k(n-3)$ .*

---

\*Institut Fourier, CNRS Université Grenoble Alpes, email: michel.mollard@univ-grenoble-alpes.fr

In [3] Elif Saygı and Ömer Eğecioğlu, solved an open problem proposed by the authors of [2]. They proved that asymptotically all vertices of  $\Gamma_n$  are covered by a maximum set of disjoint subgraphs isomorphic to  $Q_k$  thus that

**Theorem 1.2** *For every  $k \geq 1$ ,  $\lim_{n \rightarrow \infty} \frac{q_k(n)}{|V(\Gamma_n)|} = \frac{1}{2^k}$ .*

The ingenious, but long, proof they proposed is a nine cases study of the decomposition of the generating function of  $q_k(n)$ . The purpose of this short note is to deduce from Theorem 1.1 a recursive formula for the number of non covered vertices by a maximum set of disjoint hypercubes. We obtain as a consequence an immediate proof of Theorem 1.2.

## 2 Number of non covered vertices

**Definition 2.1** *Let  $\{P_k(n)\}_{k=1}^{\infty}$  be the family of sequences of integers defined by*

*(i)  $P_k(n+3) = P_k(n) + 2P_{k-1}(n+1)$  for  $k \geq 2$  and  $n \geq 0$*

*(ii)  $P_k(0) = 1, P_k(1) = 2, P_k(2) = 3$ , for  $k \geq 2$*

*(iii)  $P_1(n) = 0$  if  $n \equiv 1[3]$  and  $P_1(n) = 1$  if  $n \equiv 0[3]$  or  $n \equiv 2[3]$ .*

Solving the recursion consecutively for the first values of  $k$  and each class of  $n$  modulo 3 we obtain the first values of  $P_k(n)$ .

$n \bmod 3$	0	1	2
$P_1(n)$	1	0	1
$P_2(n)$	1	$\frac{2}{3}n + \frac{4}{3}$	$\frac{2}{3}n + \frac{5}{3}$
$P_3(n)$	$\frac{2}{9}n^2 + \frac{2}{3}n + 1$	$\frac{2}{9}n^2 + \frac{8}{9}n + \frac{8}{9}$	$\frac{2}{9}n^2 + \frac{10}{9}n + \frac{10}{9}$
$P_4(n)$	$\frac{4}{81}n^3 + \frac{2}{9}n^2 + \frac{2}{9}n + 1$	$\frac{2}{9}n^2 + \frac{8}{9}n + \frac{8}{9}$	$\frac{4}{81}n^3 + \frac{4}{27}n^2 + \frac{10}{27}n + \frac{103}{81}$

Table 1:  $P_k(n)$  for  $k = 1, \dots, 4$

**Proposition 2.2** *Let  $n = 3p+r$  with  $r = 0, 1$  or  $2$ . For a fixed  $r$ ,  $P_k(n)$  is a polynomial in  $n$  of degree at most  $k-1$ .*

**Proof.** From (i) we can write

$$P_k(n) = 2 \sum_{i=0}^{p-1} P_{k-1}(n-2-3i) + P_k(r).$$

For any integer  $d$  the classical Faulhaber's formula expresses the sum  $\sum_{m=0}^n m^d$  as a polynomial in  $n$  of degree  $d+1$ . Thus if  $Q(n)$  is a polynomial of degree at most  $d$  then  $\sum_{m=0}^n Q(m)$  is a polynomial in  $n$  of degree at most  $d+1$ . Let  $Q'(m) = Q(m)$  if

$m \equiv 0[3]$  and 0 otherwise. Applying this to  $Q'$  we obtain that  $\sum_{m=0, m \equiv 0[3]}^n Q(m)$  is also a polynomial in  $n$  of degree at most  $d+1$ . Thus if  $P_{k-1}(n)$  is a polynomial in  $n$  of degree at most  $k-2$  then  $\sum_{i=0}^{p-1} P_{k-1}(n-2-3i)$  is a polynomial of degree at most  $k-1$ . Since for a fixed  $r$   $P_1(n)$  is a constant, by induction on  $k$ ,  $P_k(n)$  is a polynomial in  $n$  of degree at most  $k-1$ .  $\square$

**Theorem 2.3** *The number of non covered vertices of  $\Gamma_n$  by  $q_k(n)$  disjoint  $Q_k$ 's is  $P_k(n)$ .*

**Proof.** This is true for  $k=1$  since the Fibonacci cube  $\Gamma_n$  has a perfect matching for  $n \equiv 1[3]$  and a maximum matching missing a vertex otherwise.

For  $k > 1$  this is true for  $n=0, 1, 2$  since the values of  $P_k(n)$  are respectively 1, 2, 3 thus are equal to  $|V(\Gamma_n)|$  and there is no  $Q_k$  in  $\Gamma_n$ .

Assume the property is true for some  $k \geq 1$  and any  $n$ . Then consider  $k+1$ . By induction on  $n$  we can assume that the property is true for  $\Gamma_{n-3}$ . Let us prove it for  $\Gamma_n$ .

From Theorem 1.1 we have  $q_{k+1}(n) = q_k(n-2) + q_{k+1}(n-3)$ .

Thus the number of non covered vertices of  $\Gamma_n$  by  $q_{k+1}(n)$  disjoint  $Q_{k+1}$ 's is

$$|V(\Gamma_n)| - 2^{k+1}q_{k+1}(n) = F_{n+2} - 2^{k+1}[q_k(n-2) + q_{k+1}(n-3)] = F_{n+2} - 2 \cdot 2^k q_k(n-2) - 2^{k+1}q_{k+1}(n-3).$$

Using equalities  $P_k(n-2) = F_n - 2^k q_k(n-2)$  and  $P_{k+1}(n-3) = F_{n-1} - 2^{k+1}q_{k+1}(n-3)$  we obtain

$$|V(\Gamma_n)| - 2^{k+1}q_{k+1}(n) = F_{n+2} + 2(P_k(n-2) - F_n) + P_{k+1}(n-3) - F_{n-1}.$$

From  $F_{n+2} - 2F_n - F_{n-1} = 0$  and  $2P_k(n-2) + P_{k+1}(n-3) = P_{k+1}(n)$  the number of non covered vertices is  $P_{k+1}(n)$ . So the theorem is proved.

$\square$

For any  $k$ , since the number of non covered vertices is polynomial in  $n$  and  $|V(\Gamma_n)| = F_{n+2} \sim \frac{3+\sqrt{5}}{2\sqrt{5}}(\frac{1+\sqrt{5}}{2})^n$  we obtain, like in [3], that

$$\lim_{n \rightarrow \infty} \frac{P_k(n)}{|V(\Gamma_n)|} = 0$$

thus

$$\lim_{n \rightarrow \infty} \frac{q_k(n)}{|V(\Gamma_n)|} = \frac{1}{2^k}$$

## References

- [1] S. Klavžar, Structure of Fibonacci cubes: a survey, J. Comb. Optim. 25 (2011) 1–18.
- [2] Sylvain Gravier, Michel Mollard, Simon Špacapan, Sara Sabrina Zemljič, On disjoint hypercubes in Fibonacci cubes, Discrete Applied Mathematics, Volumes 190–191(2015) 50-55, <http://dx.doi.org/10.1016/j.dam.2015.03.016>.
- [3] Elif Saygı, Ömer Egecioğlu, Counting Disjoint Hypercubes in Fibonacci cubes, Discrete Applied Mathematics, Volume 215 (2016) 231-237, <http://dx.doi.org/10.1016/j.dam.2016.07.004>